

## AMENDMENTS

### In the Specification

Please replace the paragraph beginning on page 15, line 9, with the following rewritten paragraph:

a' --Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length  $W$  in the query sequence, which either match or satisfy some positive-valued threshold score  $T$  when aligned with a word of the same length in a database sequence.  $T$  is referred to as the neighborhood word score threshold. These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters  $M$  (reward score for a pair of matching residues; always  $> 0$ ) and  $N$  (penalty score for mismatching residues; always  $< 0$ ). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity  $X$  from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters  $W$ ,  $T$ , and  $X$  determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength ( $W$ ) of 11, an expectation ( $E$ ) of 10, a cutoff of 100,  $M=5$ ,  $N=-4$ , and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength ( $W$ ) of 3, an expectation ( $E$ ) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915).--

Please replace the paragraph beginning on page 64, line 18, with the following rewritten paragraph:

92 -- Gene identities can be determined by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) J. Mol. Biol. 215:403-410) searches under default parameters for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences are analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm. The DNA sequences are translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish, W. and States, D. J. *Nature Genetics* 3:266-272 (1993)) provided by the NCBI. In some cases, the sequencing data from two or more clones containing overlapping segments of DNA are used to construct contiguous DNA sequences.--

93 { Please replace the paragraph beginning on page 20, line 28 as follows: }

-- Plasmids containing the nucleotide sequences of the invention were deposited with the Patent Depository of the American Type Culture Collection (ATCC), Manassas, Virginia, on October 1, 2001 and assigned Patent Deposit No. PTA-3737. The deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. The deposit was made merely as a convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112. --

94 { Please replace the paragraph beginning on page 21, line 20 as follows: }

-- (g) polynucleotides comprising the sequences obtained from the clones deposited in a bacterial host with the American Type Culture Collection (ATCC) on October 1, 2001, and assigned Accession Number PTA-3737; and --